

Singular Monopoles and Gravitational Instantons

Sergey A. Cherkis*

*California Institute of Technology
Pasadena, CA 91125*

Anton Kapustin†

*School of Natural Sciences, Institute for Advanced Study
Olden Lane, Princeton, NJ 08540*

Abstract

We model A_k and D_k asymptotically locally flat gravitational instantons on the moduli spaces of solutions of $U(2)$ Bogomolny equations with prescribed singularities. We study these moduli spaces using Ward correspondence and find their twistor description. This enables us to write down the Kähler potential for A_k and D_k gravitational instantons in a relatively explicit form.

*Research supported in part by DOE grant DE-FG03-92-ER40701

†Research supported in part by DOE grant DE-FG02-90-ER40542

1 Introduction

A gravitational instanton is a smooth four-dimensional manifold with a Riemannian metric satisfying Einstein equations. A particularly interesting class of gravitational instantons is that of four-dimensional hyperkähler manifolds, i.e. manifolds with holonomy group contained in $SU(2)$. A hyperkähler manifold can be alternatively characterized as a Riemannian manifold admitting three covariantly constant complex structures I, J, K satisfying the quaternion relations

$$IJ = -JI = K, \text{ etc.} \quad (1)$$

such that the metric is Hermitian with respect to I, J, K . Covariant constancy of I, J, K implies that three 2-forms $\omega_1 = g(I\cdot, \cdot), \omega_2 = g(J\cdot, \cdot), \omega_3 = g(K\cdot, \cdot)$ are closed. If we pick one of the complex structures, say I , we may regard a hyperkähler manifold as a complex manifold equipped with Kähler metric (with Kähler form ω_1) and a complex symplectic form $\omega = \omega_2 + i\omega_3$.

Hyperkähler four-manifolds arise in several physical problems. For example, compactification of string and M-theory on hyperkähler four-manifolds preserves one half of supersymmetries and provides exact solutions of stringy equations of motion.

The only compact hyperkähler four-manifolds are T^4 and K3, but the K3 metric is not known explicitly. In the noncompact case there are several possibilities to consider. There are no nontrivial hyperkähler metrics asymptotically approaching that of $\mathbf{R}^4$, but the situation becomes more interesting if one asks that the metric be only Asymptotically Locally Euclidean (ALE), i.e. that the metric look asymptotically like the quotient of $\mathbf{R}^4$ by a finite group of isometries. All such metrics fit into the ADE classification of Kronheimer which we now briefly explain. Let Γ be a finite subgroup of $SU(2)$. There is a natural correspondence between such Γ 's and ADE Dynkin diagrams: A_k diagram corresponds to the cyclic group $\mathbf{Z}_{k+1}$, D_k diagram corresponds to the binary dihedral group $\mathbf{D}_{k-2}$ of order $4(k-2)$, and E_k diagrams correspond to symmetry groups of tetrahedron, cube, and icosahedron. Since $SU(2)$ acts on $\mathbf{C}^2$ by the fundamental representation, we may consider quotients $\mathbf{C}^2/\Gamma$ (known as Kleinian singularities). Kronheimer showed that resolutions of Kleinian singularities admit ALE hyperkähler metrics, and that all such metrics arise in this way [1, 2]. In the A_k case the metric has been known explicitly for some time: it is the Gibbons-Hawking metric with $k+1$ centers [3]. Kronheimer provided an implicit construction of D_k and E_k ALE

gravitational instantons as hyperkähler quotients [1].

Another interesting class of noncompact gravitational instantons is that of Asymptotically Locally Flat (ALF) manifolds. This means that the metric asymptotically approaches the metric on $(\mathbf{S}^1 \times \mathbf{R}^3)/\Gamma$ where Γ is some finite group. The only known hyperkähler metric of this sort is the multi-Taub-NUT metric. As a complex manifold the $k+1$ -center multi-Taub-NUT space is isomorphic to the resolution of $\mathbf{C}^2/\mathbf{Z}_{k+1}$, so we will call it A_k -type ALF gravitational instanton. Compactification of M-theory of this manifold is equivalent to a configuration of $k+1$ parallel D6 branes in IIA string theory. Furthermore, it is expected that a configuration of an O6+ orientifold and k D6 branes in IIA string theory corresponds to the compactification of M-theory on a D_k ALF space [4], i.e. an ALF gravitational instanton isomorphic to the resolution of $\mathbf{C}^2/\mathbf{D}_{k-2}$. More generally, any compactification of M-theory on an ALF hyperkähler manifold should correspond to a IIA brane configuration preserving half of supersymmetries. Thus it is of interest to find all four-dimensional ALF hyperkähler metrics in as explicit form as possible.

In our previous paper [5] we constructed D_k ALF metrics from moduli spaces of certain ordinary differential equations (Nahm equations). In this paper we construct both A_k and D_k ALF hyperkähler four-manifolds from moduli spaces of solutions of $U(2)$ Bogomolny equations on $\mathbf{R}^3$ with prescribed singularities. Solutions of $SU(2)$ Bogomolny equations with singularities were previously considered by Kronheimer [6], and much of our discussion closely follows that in Ref. [6]. The idea is that, on one hand, the moduli space of Bogomolny equations carries natural hyperkähler structure, while on the other hand the solutions can be found by means of Ward correspondence. This approach yields directly the twistor space (in the sense of Penrose) of the moduli space of solutions. To get the metric itself one needs to find an appropriate family of sections of the twistor space. In both A_k and D_k cases we were able to identify the correct family of sections only modulo some finite choices. The A_k metrics are simple enough so that one can explicitly see that only one choice gives nonsingular metrics. In the end of subsection 5.1 we argue that the D_k twistor spaces we have constructed correspond to everywhere smooth hyperkähler metrics as well. We also show that the D_k ALF metrics we obtain are identical to those found in Ref. [5].

Let us explain what we mean by solutions of Bogomolny equations with prescribed singularities. Recall that Bogomolny equations on $\mathbf{R}^3$ are equations for a connection A in a vector bundle B over $\mathbf{R}^3$ and a section Φ of

EndB:

$$*F_A = D_A\Phi.$$

Let $\|a\|$ be the Ad-invariant norm on $u(2)$, $\|a\|^2 = -\frac{1}{2}\text{Tra}^2$. Fix k distinct points $\vec{p}_1, \dots, \vec{p}_k \in \mathbf{R}^3$. A *singular $U(2)$ monopole* is a solution of $U(2)$ Bogomolny equations on $\mathbf{R}^3 \setminus \{\vec{p}_1, \dots, \vec{p}_k\}$ satisfying the following conditions.

- (i) As $\vec{r} \rightarrow \vec{p}_\alpha$ $2r_\alpha\Phi \rightarrow i \text{diag}(0, -\ell_\alpha)$ up to gauge transformations, and $d(r_\alpha\|\Phi\|)$ is bounded. Here $r_\alpha = |\vec{r} - \vec{p}_\alpha|$, $\alpha = 1, \dots, k$.
- (ii) As $r \rightarrow \infty$ one has asymptotic expansions, up to gauge transformations,

$$\Phi = i \text{diag} \left(\mu_1 - \frac{n}{2r}, \mu_2 + \frac{n - \sum \ell_\alpha}{2r} \right) + O(1/r^2),$$

$$\frac{\partial\|\Phi\|}{\partial\Omega} = O(1/r^2), \quad \|D\Phi\| = O(1/r^2).$$

We will refer to n as the nonabelian charge of the monopole, and to $\{\ell_\alpha\}$ as its abelian charges. We will assume that $\mu_1 > \mu_2$. We also set $n' = n - \sum_\alpha \ell_\alpha$, $\mu = \mu_1 - \mu_2$, for short.

Every fiber of the complex rank two bundle B splits into the eigenspaces of Φ , $B = M_1 \oplus M_2$, near $\vec{r} = \vec{p}_\alpha$ or when $r \rightarrow \infty$. Let M_1 correspond the eigenvalue of Φ diverging as $\vec{r} \rightarrow \vec{p}_\alpha$. It is a simple consequence of Bogomolny equations that $-\ell_\alpha$ is the degree of M_1 restricted to a small 2-sphere around $\vec{r} = \vec{p}_\alpha$. Similarly, $-n$ and n' are the degrees of eigensubbundles of B restricted to a large 2-sphere. Therefore n and $\{\ell_\alpha\}$ are integers.

String theory considerations imply that the moduli space of $n = 1$ monopole with $\ell_\alpha = 1, \alpha = 1, \dots, k$ is an A_{k-1} ALF gravitational instanton, and the centered moduli space of $n = 2$ monopole¹ with $\ell_\alpha = 1, \alpha = 1, \dots, k$ is a D_k ALF gravitational instanton [7]. In this paper we show that this is indeed the case. The main tool is the Ward correspondence described in section 2. In section 3 we use it to derive the twistor space for arbitrary n . In sections 4 and 5 we deal with $n = 1$ and $n = 2$ cases, respectively. We show that the moduli spaces are resolutions of A_{k-1} and D_k singularities, as expected, find the real holomorphic sections of the twistor spaces, and derive the Kähler potentials for the metrics using the generalized Legendre transform method of Refs. [8, 9].

¹The centered $n = 2$ monopole moduli space is a $U(1)$ hyperkähler quotient of the $n = 2$ monopole moduli space; see section 5 below.

2 Ward correspondence

From now on we restrict ourselves to the case $\ell_\alpha = 1, \alpha = 1, \dots, k$. We make some comments on the more general case of positive ℓ_α at the end of subsection 5.1. To construct the moduli space of singular $U(2)$ monopoles, we will use a version of Ward correspondence due to Hitchin [10]. The set of all oriented straight lines $\mathbf{T}$ in $\mathbf{R}^3$ has a natural complex structure, as it is the tangent bundle of the projective line. $\mathbf{T}$ can be covered by two patches $V_0(\zeta \neq \infty)$ and $V_1(\zeta \neq 0)$ with coordinates (η, ζ) and $(\eta', \zeta') = (\eta/\zeta^2, 1/\zeta)$. For any point $\vec{x} \in \mathbf{R}^3$ the set of all oriented straight lines through $\vec{x}$ sweeps out a projective line $P_x \in \mathbf{T}$; thus there is a holomorphic map $P_x : \mathbf{P}^1 \rightarrow \mathbf{T}$. The reversal of the orientation of lines in $\mathbf{R}^3$ is an antiholomorphic map $\tau : \mathbf{T} \rightarrow \mathbf{T}$ satisfying $\tau^2 = id$. It is called the real structure of $\mathbf{T}$. For any $\vec{x}$ it acts on P_x as the antipodal map. Thus P_x is a real holomorphic section of $\mathbf{T}$.

For any straight line in $\mathbf{R}^3$

$$\gamma = \{\vec{x} | \vec{x} = \vec{u}t + \vec{v}, \vec{u} \cdot \vec{u} = 1, \vec{u} \cdot \vec{v} = 0\}$$

let

$$\begin{aligned} \gamma_+ &= \{\vec{x} | \vec{x} = \vec{u}t + \vec{v}, t > R\}, \\ \gamma_- &= \{\vec{x} | \vec{x} = \vec{u}t + \vec{v}, t < -R\}, \end{aligned} \tag{2}$$

where R is a positive number greater than any $|\vec{p}_\alpha|$. Now we define two complex rank 2 vector bundles E^+ and E^- over $\mathbf{T}$:

$$\begin{aligned} E^+ &= \{s \in \Gamma(\gamma_+, E) | D_\gamma s = i\Phi s\}, \\ E^- &= \{s \in \Gamma(\gamma_-, E) | D_\gamma s = i\Phi s\}. \end{aligned} \tag{3}$$

From Bogomolny equations it follows, as in Ref. [10], that these bundles are holomorphic. The real structure τ on $\mathbf{T}$ can be lifted to an antilinear antiholomorphic map

$$\sigma : E^+ \rightarrow (E^-)^*.$$

Thus every solution of $U(2)$ Bogomolny equations maps to a pair of holomorphic rank two bundles on $\mathbf{T}$ interchanged by the real structure.

Let P_x denote the real section corresponding to $\vec{x}$, and P_α the real section corresponding to $\vec{p}_\alpha$. Let P be the union of all P_α . If γ does not pass through any of $\vec{p}_\alpha$, any solution s can be continued from γ_+ to γ_- . This

defines a natural identification of the fibers E_γ^+ and E_γ^- . Therefore we have an isomorphism

$$h : E^+|_{\mathbf{T} \setminus P} \rightarrow E^-|_{\mathbf{T} \setminus P}. \quad (4)$$

For nonsingular monopoles h extends to an isomorphism over the whole $\mathbf{T}$, therefore the Ward correspondence maps a nonsingular monopole to a holomorphic bundle over $\mathbf{T}$. In the present case h or h^{-1} may have singularities at P , and the Ward correspondence maps a singular monopole into a triplet (E^+, E^-, h) . This triplet satisfies a certain triviality constraint which we now proceed to formulate.

For any $\vec{x}$ distinct from all $\vec{p}_\alpha$ the intersection $P_x \cap P$ consists of an even number of points. For a generic $\vec{x}$ the cardinality of $Q_x = P_x \cap P$ is $2k$. For any $\vec{x}$ we can arbitrarily split Q_x into two sets of equal cardinality Q_x^+ and Q_x^- and construct a vector bundle E_x over P_x by gluing together E^+ restricted to $P_x \setminus Q_x^+$ and E^- restricted to $P_x \setminus Q_x^-$, with the transition function h . (Of course, E_x depends on the splitting.) The triviality constraint is that for any $\vec{x}$ there is a splitting $Q_x = Q_x^+ \cup Q_x^-$ such that E_x is trivial.

Now we state the Ward correspondence between singular $U(2)$ monopoles and twistor data. There is a bijection² between singular monopoles modulo gauge transformations and pairs (E^+, E^-) of holomorphic rank 2 bundles over $\mathbf{T}$ equipped with an isomorphism Eq. (4) satisfying the following conditions:

(a) For any $\vec{x} \neq \vec{p}_\alpha$ there is a splitting $Q_x = Q_x^+ \cup Q_x^-$ such that E_x is trivial.

(b) In the vicinity of each point of P there exist trivializations of E^+ and E^- such that h takes the form

$$h = \begin{pmatrix} 1 & 0 \\ 0 & \Pi_\alpha(\eta - P_\alpha(\zeta)) \end{pmatrix}, \quad (5)$$

so that h extends across P to a morphism $E^+ \rightarrow E^-$.

(c) The real structure τ on $\mathbf{T}$ lifts to an antilinear antiholomorphic map $\sigma : E^+ \rightarrow (E^-)^*$.

Let us explain where (a) and (b) come from. The condition (b) arises from studying the behavior of the solutions of the equation $D_\gamma s = i\Phi s$ as γ approaches $\vec{p}_\alpha$. Details can be found in Ref. [6]. (There the $SU(2)$ case was analyzed, but the extension to $U(2)$ is straightforward). To demonstrate

²The injectivity of Ward correspondence can be shown by a straightforward modification of the argument in Ref. [10]. We conjecture the surjectivity by analogy with the nonsingular case.

(a) it is sufficient to exhibit a holomorphic trivialization of E_x . Take any $\vec{x} \neq \vec{p}_\alpha, \alpha = 1, \dots, k$ and recall that P_x consists of all straight lines γ passing through $\vec{x}$. To obtain a holomorphic section of E_x pick a vector v_1 in the fiber of B over $\vec{x}$ and take it as an initial condition for the equation $D_\gamma s = i\Phi s$ at $t = 0$. Integrating it forward and backward in t and varying γ yields sections of E^+ and E^- related by h . It is easy to check that they are holomorphic and thereby combine into a holomorphic section s_1 of E_x . To get a section s_2 of E_x linearly independent from s_1 just pick a vector v_2 linearly independent from v_1 and repeat the procedure. (This argument has to be modified if there is a straight line γ passing through $\vec{x}$ and $\alpha, \beta \in \{1, \dots, k\}$ such that $\vec{p}_\alpha$ and $\vec{p}_\beta$ lie on γ and $\vec{x}$ separates them. In this case one of the vectors v_1, v_2 has to be varied, $v_i \sim \zeta^{-1}$, as one varies γ .)

We now want to encode the twistor data in an algebraic curve $S \subset \mathbf{T}$, in the spirit of Ref. [10]. We denote by $\mathcal{O}(m)$ the pullback to $\mathbf{T}$ of the unique degree m line bundle on $\mathbf{P}^1$, and by $L^x(m)$ a line bundle over $\mathbf{T}$ with the transition function $\zeta^{-m} e^{-x\eta/\zeta}$ from V_0 to V_1 . Let L_1^+ be a line subbundle of E^+ which consists of solutions of $D_\gamma s = i\Phi s$ bounded by $\text{const} \cdot \exp(-\mu_1 t)t^n$ as $t \rightarrow +\infty$. Similarly, a line bundle $L_1^- \subset E^-$ consists of solutions bounded by $\text{const} \cdot \exp(-\mu_2 t)t^{-n'}$ as $t \rightarrow -\infty$. The line bundles L_2^+ and L_2^- are defined by

$$L_2^+ = E^+ / L_1^+, \quad L_2^- = E^- / L_1^-.$$

As in Ref. [10] the asymptotic conditions on the Higgs field can be used to show that $L_{1,2}^+$ and $L_{1,2}^-$ are holomorphic line bundles, and that the following isomorphisms hold:

$$L_1^+ \simeq L^{\mu_1}(-n), \quad L_2^+ \simeq L^{\mu_2}(n'), \quad L_1^- \simeq L^{\mu_2}(-n'), \quad L_2^- \simeq L^{\mu_1}(n).$$

Consider a composite map

$$\psi : L_1^+ \rightarrow E^+ \rightarrow E^- \rightarrow L_2^-,$$

where the first arrow is an inclusion, the second arrow is h , and the third arrow is a natural projection. We may regard ψ as an element of $H^0(\mathbf{T}, \mathcal{O}(2n))$. Let us define the spectral curve S to be the zero level of ψ . S is in the linear system $\mathcal{O}(2n)$. Arguments identical to those in Ref. [10] can be used to prove that S is compact and real (i.e., $\tau(S) = S$).

Consider now a map $\phi : \wedge^2 E^+ \rightarrow \wedge^2 E^-$ induced by h . By virtue of Eq. (5) the zero level of ϕ is precisely P . We will assume in what follows that S does

not contain any of P_α as components. Physically this corresponds to the requirement that none of the nonabelian monopoles is located at $\vec{x} = \vec{p}_\alpha$. For simplicity we will also assume that $S \cap P$ consists of $2nk$ points (this is a generic situation).

The construction here bears a close resemblance to that in Ref. [11], where nonsingular monopoles for all classical groups were constructed. According to Ref. [11], the spectral data for a nonsingular $SU(3)$ monopole with magnetic charge (k, n) include a pair of spectral curves S_1, S_2 in the linear systems $\mathcal{O}(2n), \mathcal{O}(2k)$. Our S and P are analogs of S_1 and S_2 . The condition that $S \cap P$ consists of $2nk$ points is analogous to the requirement in Ref. [11] that the monopoles are generic. (This resemblance is not a coincidence: if we consider an $SU(3)$ gauge theory broken down to $SU(2) \times U(1)$ by a large vev of an adjoint Higgs field, the (k, n) monopoles of $SU(3)$ reduce to singular monopoles of $SU(2) \times U(1)$ with nonabelian charge n and total abelian charge k . In this limit the spectral data of Ref. [11] must reduce to ours.)

Since $L_1^+|_S = \ker \psi|_S$, we have a well-defined holomorphic map $\rho : L_2^+|_S \rightarrow L_2^-|_S$ induced by h . There is also a holomorphic map $\xi : L_1^+|_S \rightarrow L_1^-|_S$ induced by h . Thus we have natural elements $\rho \in H^0(S, L^\mu(k))$ and $\xi \in H^0(S, L^{-\mu}(k))$. It also easily follows from the definition that $\rho \otimes \xi = \phi|_S$, and therefore the divisors of both ρ and ξ are subsets of $S \cap P$. ρ and ξ are interchanged by real structure, and therefore the same is true about their divisors. It follows that the divisors of ρ and ξ are disjoint and have equal cardinality. Thus we can define the spectral data for a generic singular monopole to consist of

- (i) A spectral curve S , which is a real compact curve in the linear system $\mathcal{O}(2n)$ such that $S \cap P$ consists of $2nk$ disjoint points.
- (ii) A splitting $S \cap P = Q^+ \cup Q^-$ into sets of equal cardinality interchanged by τ .
- (iii) A section ρ of $L^\mu(k)|_S$ with divisor Q^+ and a section ξ of $L^{-\mu}(k)|_S$ with divisor Q^- . ρ and ξ are interchanged by real structure.

The condition (iii) is a constraint on S . It implies that ρ and ξ satisfy

$$\rho\xi = \prod_{\alpha}(\eta - P_{\alpha}(\zeta)). \quad (6)$$

For nonsingular monopoles it reduces to the requirement that $L^\mu|_S$ be trivial, as in Ref. [10]. As a consequence of (iii), $L^{2\mu}|_S[Q_- - Q_+]$ is trivial.

Recall that the spectral data for nonsingular $SU(2)$ monopoles satisfy an additional constraint, the “vanishing theorem” of Ref. [12]. It says that

$L^{z\mu}(n-2)$ is nontrivial for $z \in (0, 1)$. A natural guess for the analogue of this condition in our case is

(iv) $L^{z\mu}(n-2)[-Q_+]$ is nontrivial for $z \in (0, 1)$.

We already mentioned a close connection of the spectral data for singular $U(2)$ monopoles and those for nonsingular $SU(3)$ monopoles [11] with the largest Higgs vev set to $+\infty$. Consequently, one can obtain the condition (iv) from the “vanishing theorem” of Ref. [11] by taking the appropriate limit. A direct derivation of (iv) should also be possible.

Arguments very similar to those in Ref. [10] show that the spectral data determine the singular monopole uniquely. A natural question is if there is a one-to-one correspondence between singular $U(2)$ monopoles and spectral data defined by (i-iv). The answer was positive for nonsingular $SU(2)$ monopoles [12], so it is highly plausible that the same is true in the present case. Presumably a proper proof of this can be achieved by converting the spectral data into solutions of Nahm equations, and then reconstructing singular monopoles by an inverse Nahm transform [12, 11].

3 Twistor space for singular monopoles

Having established the correspondence between singular $U(2)$ monopoles and algebraic data on $\mathbf{T}$, we now proceed to construct the twistor space Z_n for the moduli space of a singular monopole with nonabelian charge n . We follow the method of Ref. [13]. For fixed $\zeta = \zeta_0$ every point in Z_n yields a spectral curve S which intersects the fiber of $\mathbf{T}$ over ζ_0 at n points. Thus we have a projection

$$Z_n \rightarrow \oplus_{j=1}^n \mathcal{O}(2j) = Y_n.$$

Concretely, if S is given by $\eta^n + \eta_1\eta^{n-1} + \dots + \eta_n = 0$, the corresponding point in Y_n is $(\eta_1, \dots, \eta_n)$. Now consider an n -fold cover of Y_n

$$X_n = \left\{ (\eta, \eta_1, \dots, \eta_n) \in \mathcal{O}(2) \oplus Y_n \mid \eta^n + \eta_1\eta^{n-1} + \dots + \eta_n = 0 \right\}.$$

There are two natural projections $\pi_1 : X_n \rightarrow \mathbf{T}$ and $\pi_2 : X_n \rightarrow Y_n$. Using these projections, we get a rank n bundle V^+ over Y_n as a direct image sheaf $V^+ = \pi_{2*}\pi_1^*L^\mu(k)$. Similarly, we get a rank n bundle $V^- = \pi_{2*}\pi_1^*L^{-\mu}(k)$. For any point in Z_n we have a section ρ of $L^\mu(k)|_S$ and a section ξ of $L^{-\mu}(k)|_S$. Therefore, there is an inclusion $Z_n \subset V^+ \oplus V^-$. To describe this inclusion more concretely, we must rewrite the condition (iii) in terms of sections of

$V^\pm$. The result is as follows. Let U be a $2n+1$ -dimensional subvariety in $\mathbf{C}^{3n+1}$ with coordinates $(\zeta, \eta_1, \dots, \eta_n, \rho_0, \dots, \rho_{n-1}, \xi_0, \dots, \xi_{n-1})$ defined by

$$\begin{aligned} (\rho_0 + \rho_1\eta + \dots + \rho_{n-1}\eta^{n-1})(\xi_0 + \xi_1\eta + \dots + \xi_{n-1}\eta^{n-1}) &= \prod_{\alpha} (\eta - P_{\alpha}(\zeta)), \\ \text{mod } \eta^n + \eta_1\eta^{n-1} + \dots + \eta_n &= 0. \end{aligned} \quad (7)$$

Take two copies of U and glue them together over $\zeta \neq 0, \infty$ by

$$\begin{aligned} \tilde{\zeta} &= \zeta^{-1}, \\ \tilde{\eta}_j &= \zeta^{-2j}\eta_j, \quad j = 1, \dots, n, \\ \tilde{\rho}_0 + \tilde{\rho}_1\tilde{\eta} + \dots + \tilde{\rho}_{n-1}\tilde{\eta}^{n-1} &= e^{-\mu\eta/\zeta}\zeta^{-k}(\rho_0 + \rho_1\eta + \dots + \rho_{n-1}\eta^{n-1}), \\ \tilde{\xi}_0 + \tilde{\xi}_1\tilde{\eta} + \dots + \tilde{\xi}_{n-1}\tilde{\eta}^{n-1} &= e^{\mu\eta/\zeta}\zeta^{-k}(\xi_0 + \xi_1\eta + \dots + \xi_{n-1}\eta^{n-1}), \end{aligned} \quad (8)$$

all modulo $\eta^n + \eta_1\eta^{n-1} + \dots + \eta_n = 0$. The resulting $2n+1$ -dimensional variety is Z_n , the twistor space of singular monopoles with nonabelian charge n .

To reconstruct the hyperkähler metric from the twistor space one has to find a holomorphic section of $\Lambda^2 T_F^* \otimes \mathcal{O}(2)$, where T_F^* is the cotangent bundle of the fiber of Z_n . Upon restriction to any fiber of Z_n this section must be closed and nondegenerate. An obvious choice (the same as in Ref. [13]) is

$$\omega = 4 \sum_{j=1}^n \frac{d\rho(\beta_j) \wedge d\beta_j}{\rho(\beta_j)}, \quad (9)$$

where $\beta_j, j = 1, \dots, n$ are the roots of $\eta^n + \eta_1\eta^{n-1} + \dots + \eta_n = 0$.

4 Moduli space M_1 of $n = 1$ monopole

Specializing the formulas of the previous section to $n = 1$, we get that the twistor space Z_1 is a hypersurface in the total space of $L^\mu(k) \oplus L^{-\mu}(k)$

$$\rho_0 \xi_0 = \prod_{\alpha=1}^k (\eta - P_{\alpha}(\zeta)), \quad (10)$$

where $\rho_0 \in L^\mu(k)$, $\xi_0 \in L^{-\mu}(k)$, and $\eta \in \mathcal{O}(2)$. Obviously, for fixed ζ this is a resolution of $\mathbf{C}^2/\mathbf{Z}_k$, so the corresponding hyperkähler metric is an A_{k-1} gravitational instanton. In fact, it is well known what the metric is: it is the multi-Taub-NUT metric with k centers. In the remainder of this section we

rederive this result using the Legendre transform method of Refs. [14, 8, 9]. This will serve as a warm-up for the discussion of D_k ALF metrics in the next section.

First we find the real holomorphic sections of the twistor space Z_1 . This amounts to solving Eq. (10) with ρ_0, ξ_0 , and η now regarded as holomorphic sections of the appropriate bundles. Recalling that $\eta = a\zeta^2 + 2b\zeta - \bar{a}$ and $P_\alpha(\zeta) = a_\alpha\zeta^2 + 2b_\alpha\zeta - \bar{a}_\alpha$ with $b, b_\alpha \in \mathbf{R}$, one gets in the patch V_0

$$\begin{aligned}\rho_0 &= Ae^{+\mu(b+a\zeta)} \prod_{\alpha=1}^k (\zeta - u_\alpha), \\ \xi_0 &= Be^{-\mu(b+a\zeta)} \prod_{\alpha=1}^k (\zeta - v_\alpha),\end{aligned}$$

with $AB = \prod (a - a_\alpha)$. Here u_α and v_α are the roots of the equation $\eta(\zeta) = P_\alpha(\zeta)$,

$$\begin{aligned}u_\alpha &= \frac{-(b - b_\alpha) - \Delta_\alpha}{a - a_\alpha}, \\ v_\alpha &= \frac{-(b - b_\alpha) + \Delta_\alpha}{a - a_\alpha},\end{aligned}\tag{11}$$

with $\Delta_\alpha = \sqrt{(b - b_\alpha)^2 + |a - a_\alpha|^2} > 0$. (The ambiguity in the sign of Δ_α is fixed by requiring that the hyperkähler metric on this family of sections be everywhere nonsingular. This is equivalent to asking that the normal bundle of every section in the family be $\mathcal{O}(1) \oplus \mathcal{O}(1)$.) Since the real structure must interchange ρ_0 and ξ_0 , we get

$$B\bar{B} = \prod (b - b_\alpha + \Delta_\alpha).\tag{12}$$

Thus we have a family of solutions to Eq. (10) parametrized by $\text{Re } a$, $\text{Im } a$, b , and $\text{Arg } B$.

Having found the real holomorphic sections, we compute the Kähler potential. The twistor space Z_1 is fibered over $\mathbf{P}^1$ with an intermediate projection

$$Z_1 \rightarrow \mathcal{O}(2) \rightarrow \mathbf{P}^1.$$

In the above ζ and η are coordinates on the base and the fiber of $\mathcal{O}(2)$, respectively. The holomorphic 2-form $\omega \in \Lambda^2 T^* \otimes \mathcal{O}(2)$ is given by

$$\omega = 4d\eta \wedge \frac{d\rho}{\rho}.\tag{13}$$




Figure 1: The contour Γ_α enclosing u_α and v_α .

For $\zeta \neq \infty$ we can choose $\eta(\zeta)$ and $\chi = 2 \log \frac{\rho}{\xi}$ as two coordinates on the moduli space M_1 holomorphic with respect to the complex structure defined by ζ . The coordinates in the patch $\zeta \neq 0$ are related to these as

$$\eta' = \eta/\zeta^2, \chi' = \chi - 4\mu\eta/\zeta. \quad (14)$$

The second equation here follows from ρ_0 and ξ_0 being sections of $L^\mu(k)$ and $L^{-\mu}(k)$. In terms of these coordinates

$$\omega = d\eta \wedge d\chi = \zeta^2 d\eta' \wedge d\chi'. \quad (15)$$

Following Ref. [9] we define an auxiliary function $\hat{f}$ and a contour C by the equation

$$\oint_C \frac{d\zeta}{\zeta^j} \hat{f} = \oint_0 \frac{d\zeta}{\zeta^j} \chi + \oint_\infty \frac{d\zeta}{\zeta^j} \chi' = \left(\oint_0 + \oint_\infty \right) \frac{d\zeta}{\zeta^j} \chi - 4\mu \oint_\infty \frac{d\zeta}{\zeta^{j+1}} \eta. \quad (16)$$

for any integer j . Here and in what follows the integrals $\oint_0$ and $\oint_\infty$ are taken along small positively oriented contours around respective points. This implies that in the first of this integrals the contour runs counterclockwise, while in the second one it runs clockwise. Substituting an explicit expression for χ we find

$$\oint_C \frac{d\zeta}{\zeta^j} \hat{f} = \sum_\alpha \oint_{\Gamma_\alpha} \frac{d\zeta}{\zeta^j} 2 \log(\eta(\zeta) - P_\alpha(\zeta)) + 4\mu \oint_0 \frac{d\zeta}{\zeta^{j+1}} \eta. \quad (17)$$

Here Γ_α is a figure-eight-shaped contour enclosing u_α and v_α (see Figure 1). We define a function $G(\eta, \zeta)$ by $\partial G / \partial \eta = \hat{f}$. According to Ref. [9] the Legendre transform of the Kähler potential is given by

$$F(a, b) = \frac{1}{2\pi i} \oint_C \frac{d\zeta}{\zeta^2} G(\eta, \zeta). \quad (18)$$

Using Eq. (17) we find

$$F(a, b) = \frac{2\mu}{2\pi i} \oint_0 \frac{d\zeta}{\zeta^3} \eta^2 + \sum_{\alpha=1}^k \frac{1}{2\pi i} \oint_{\Gamma_\alpha} \frac{d\zeta}{\zeta^2} 2(\eta - P_\alpha) \log(\eta - P_\alpha). \quad (19)$$

The Kähler potential K is the Legendre transform of F :

$$K(a, \bar{a}, t, \bar{t}) = F - b(t + \bar{t}), \quad \frac{\partial F}{\partial b} = t + \bar{t}. \quad (20)$$

It is a well known fact that the metric corresponding to Eq. (19) is the multi-Taub-NUT metric with k centers [9, 14]. This is in agreement with string theory predictions [7].

5 Moduli space of centered $n = 2$ monopole

5.1 Twistor space Z_2^0 of centered $n = 2$ monopole

For $n = 2$ the moduli space M_2 is 8-dimensional and admits a triholomorphic $U(1)$ action. We define the centered moduli space M_2^0 to be the hyperkähler quotient of M_2 with respect to this $U(1)$ (at zero level). The $U(1)$ action on M_2 lifts to a $\mathbf{C}^*$ action on Z_2 . It acts by $\rho_j \rightarrow \lambda \rho_j$, $\xi_j \rightarrow \lambda^{-1} \xi_j$. The corresponding moment map is η_1 , as can be easily seen from the expression for ω . Thus Z_2^0 , the twistor space of M_2^0 , is the $\mathbf{C}^*$ quotient of the subvariety $\eta_1 = 0 = \tilde{\eta}_1$ in Z_2 . We first investigate one coordinate patch of Z_2^0 . Let us denote $\psi_1 = \rho_0 \xi_0$, $\psi_2 = \rho_1 \xi_1$, $\psi_3 = \frac{1}{2}(\rho_0 \xi_1 + \rho_1 \xi_0)$, $\psi_4 = \frac{1}{2}(\rho_0 \xi_1 - \rho_1 \xi_0)$. The variables ψ_i are invariant with respect to $\mathbf{C}^*$ action and satisfy

$$\begin{aligned} \psi_1 \psi_2 &= \psi_3^2 - \psi_4^2, \\ \psi_1 - \eta_2 \psi_2 + 2\sqrt{-\eta_2} \psi_3 &= \prod_{\alpha} (\sqrt{-\eta_2} - P_{\alpha}(\zeta)), \\ \psi_1 - \eta_2 \psi_2 - 2\sqrt{-\eta_2} \psi_3 &= \prod_{\alpha} (-\sqrt{-\eta_2} - P_{\alpha}(\zeta)). \end{aligned} \quad (21)$$

These equations define a three-dimensional subvariety U^0 in $\mathbf{C}^6$ with coordinates $(\zeta, \eta_2, \psi_1, \dots, \psi_4)$. Geometric invariant theory tells us that Z_2^0 can be obtained by gluing together two copies of U^0 over $\zeta \neq 0, \infty$. The transition functions can be computed from Eq. (8):

$$\tilde{\zeta} = \zeta^{-1}, \quad (22)$$

$$\begin{aligned}
\tilde{\eta}_2 &= \zeta^{-4}\eta_2, \\
\tilde{\psi}_1 &= \frac{\zeta^{-2k}}{2} (\psi_1 - \eta_2\psi_2 + \cos\gamma(\psi_1 + \eta_2\psi_2) - 2\psi_4\sqrt{\eta_2}\sin\gamma), \\
\tilde{\psi}_2 &= \frac{\zeta^{4-2k}}{2\eta_2} (-(\psi_1 - \eta_2\psi_2) + \cos\gamma(\psi_1 + \eta_2\psi_2) - 2\psi_4\sqrt{\eta_2}\sin\gamma), \\
\tilde{\psi}_3 &= \zeta^{2-2k}\psi_3, \\
\tilde{\psi}_4 &= \frac{\zeta^{2-2k}}{2} \left((\psi_1 + \eta_2\psi_2)\frac{\sin\gamma}{\sqrt{\eta_2}} + \psi_4\cos\gamma \right),
\end{aligned}$$

where $\gamma = 2\mu\sqrt{\eta_2}/\zeta$.

From this explicit description of Z_2^0 one can see that for any ζ the fiber of Z_2^0 is a resolution of the D_k singularity. Indeed, combining Eqs. (21) we see that the fiber of U_0 over ζ is biholomorphic to a hypersurface in $\mathbf{C}^3$ (with coordinates (η_2, ψ_2, ψ_4)) given by

$$\psi_4^2 + \eta_2\psi_2^2 + \psi_2\mathcal{Q}(\eta_2) - \mathcal{R}(\eta_2)^2 = 0, \quad (23)$$

where $\mathcal{Q}(\eta_2), \mathcal{R}(\eta_2)$ are polynomials in η_2 defined by

$$\begin{aligned}
2\mathcal{Q}(\eta_2) &= \prod_{\alpha}(\sqrt{-\eta_2} - P_{\alpha}(\zeta)) + \prod_{\alpha}(-\sqrt{-\eta_2} - P_{\alpha}(\zeta)), \\
4\sqrt{-\eta_2}\mathcal{R}(\eta_2) &= \prod_{\alpha}(\sqrt{-\eta_2} - P_{\alpha}(\zeta)) - \prod_{\alpha}(-\sqrt{-\eta_2} - P_{\alpha}(\zeta)).
\end{aligned}$$

Furthermore, these formulas imply that if all points $\vec{p}_1, \dots, \vec{p}_k$ are distinct, the manifold M_2^0 is a smooth complex manifold in any of its complex structures. Since the 2-form ω is smooth as well, we conclude that M_2^0 is a smooth hyperkähler manifold. The smoothness of M_2^0 is also in agreement with string theory predictions. Indeed, as explained in Ref. [7], the space M_2^0 is the Coulomb branch of $N = 4, D = 3$ $SU(2)$ gauge theory with k fundamental hypermultiplets, with $\vec{p}_{\alpha}$ being hypermultiplet masses. When $\vec{p}_{\alpha}$ are all distinct, the theory has no Higgs branch, and therefore the Coulomb branch is smooth everywhere. When some masses become equal, the Higgs branch emerges, and the Coulomb branch develops an orbifold singularity at the point where it meets the Higgs branch. Thus we expect that when some of $\vec{p}_{\alpha}$ coincide, or equivalently, when some of ℓ_{α} are bigger than 1, the manifold M_2^0 has orbifold singularities.

In Ref. [5] the same manifold Z_2^0 arose as the twistor space of the moduli space of a system of ordinary differential equations (so called Nahm equations). This is of course a consequence of a general correspondence between

solutions of Bogomolny equations and Nahm equations [15, 12, 11]. Thus Ref. [5] provides an equivalent construction of D_k ALF metrics.

5.2 Real holomorphic sections of Z_2^0

The discussion of section 2 implies that a real holomorphic section of the uncentered twistor space Z_2 is a triplet (S, ρ, ξ) , where S is the spectral curve in $\mathbf{T}$ given by $\eta^2 + \eta_1\eta + \eta_2 = 0$, ρ and ξ are holomorphic sections of $L^\mu(k)|_S$ and $L^{-\mu}(k)|_S$ satisfying the condition (iii) of section 2. Then, as explained in section 3, the real holomorphic sections of Z_2^0 are obtained by setting $\eta_1 = 0$ and modding by the $\mathbf{C}^*$ action $\rho \rightarrow \lambda\rho, \xi \rightarrow \lambda^{-1}\xi$. In this subsection we find the explicit form of the real holomorphic sections of Z_2^0 .

The curve $\eta^2 + \eta_2 = 0$ is either elliptic or a union of two $\mathbf{CP}^1$'s. The former case is generic, while the latter occurs at a submanifold of the moduli space. Intuitively the latter case corresponds to the situation when the two nonabelian monopoles are on top of each other. It suffices to consider the elliptic case.

By an $SO(3)$ rotation

$$\zeta = \frac{a\tilde{\zeta} + b}{-\bar{b}\tilde{\zeta} + \bar{a}}, \quad \eta = \frac{\tilde{\eta}}{(-\bar{b}\tilde{\zeta} + \bar{a})^2}, \quad |a|^2 + |b|^2 = 1, \quad (24)$$

we can always bring the elliptic curve $\eta^2 = -\eta_2(\zeta)$ to the form

$$\tilde{\eta}^2 = 4k_1^2 (\tilde{\zeta}^3 - 3k_2\tilde{\zeta}^2 - \tilde{\zeta}), \quad k_1 > 0, k_2 \in \mathbf{R}. \quad (25)$$

It follows that the discriminant $\Delta > 0$, and therefore the lattice defined by the curve S is rectangular. We denote this lattice 2Ω and its real and imaginary periods by 2ω and $2\omega'$, respectively.

We parametrized S by five real parameters: the Euler angles of the $SO(3)$ rotation and a pair of real numbers k_1 and k_2 . We will see in a moment that the condition (iii) imposes one real constraint on them, so we will obtain a four-parameter family of real sections, as required.

To write explicitly a section of $L^\mu(k)|_S$, we will use the standard “flat” parameter on the elliptic curve u defined modulo 2Ω , in terms of which $\tilde{\eta} = k_1\mathcal{P}'(u)$, $\tilde{\zeta} = \mathcal{P}(u) + k_2$. Here $\mathcal{P}(u)$ is the Weierstrass elliptic function. In terms of u the real structure acts by $u \rightarrow -\bar{u} + \omega + \omega'$.

A section of $L^\mu(k)|_S$ can be thought of as a pair of functions on S f_1, f_2 such that f_1 is holomorphic everywhere except $\zeta = \infty$, f_2 is holomorphic

everywhere except $\zeta = 0$, and for $\zeta \neq 0, \infty$ $f_2(\zeta) = \zeta^{-k} \exp(-\mu\eta/\zeta) f_1(\zeta)$. The point $\zeta = \infty$ corresponds to two points $u_\infty, -u_\infty$ on S defined by $\mathcal{P}(u_\infty) + k_2 = \bar{a}/\bar{b}$. Furthermore, condition (iii) implies that the divisor of f_1 is Q_+ . Let us recall that $Q_+ \cup Q_- = \bigcup_\alpha Q_\alpha$, where $Q_\alpha = S \cap P_\alpha, \alpha = 1, \dots, k$. Thus Q_α consists of solutions of a system of two equations $\eta = P_\alpha(\zeta), \eta^2 = -\eta_2(\zeta)$. Obviously, this defines four points on the elliptic curve S . Because of real structure, these four points split into two pairs whose members are interchanged by τ . Q_+ includes one point from each pair (for all α), Q_- includes the rest.³ Let us denote the “flat” coordinates of points in Q_+ by $u_\alpha, u'_\alpha, \alpha = 1, \dots, k$, and those in Q_- by $v_\alpha, v'_\alpha, \alpha = 1, \dots, k$. By definition, $v_\alpha = -\bar{u}_\alpha + \omega + \omega' \pmod{2\Omega}, v'_\alpha = -\bar{u}'_\alpha + \omega + \omega' \pmod{2\Omega}$. We fix the mod 2Ω ambiguity by requiring that $u_\alpha, u'_\alpha, v_\alpha, v'_\alpha$ be in the fundamental rectangle of 2Ω . In this notation a section of $L^\mu(k)|_S$ is given by

$$f_1 \sim \exp(-\mu k_1(\zeta_W(u + u_\infty) + \zeta_W(u - u_\infty)) + Cu) \prod_\alpha \frac{\sigma(u - u_\alpha)\sigma(u - u'_\alpha)}{\sigma(u - u_\infty)\sigma(u + u_\infty)}. \quad (26)$$

Here $\zeta_W(u)$ and $\sigma(u)$ are Weierstrass quasielliptic functions (we denote Weierstrass ζ -function by $\zeta_W(u)$ to avoid confusion with the affine coordinate ζ on the $\mathbf{P}^1$ of complex structures), and C is a constant. Similarly, a section of $L^{-\mu}(k)|_S$ with the divisor Q_- is represented by a pair of functions g_1, g_2 related by $g_2(\zeta) = \zeta^{-k} \exp(\mu\eta/\zeta) g_1(\zeta)$. Explicitly g_1 is given by

$$g_1 \sim \exp(\mu k_1(\zeta_W(u + u_\infty) + \zeta_W(u - u_\infty)) + Du) \prod_\alpha \frac{\sigma(u - v_\alpha)\sigma(u - v'_\alpha)}{\sigma(u - u_\infty)\sigma(u + u_\infty)}, \quad (27)$$

where D is another constant. In general f_1 and g_1 are quasiperiodic with periods 2ω and $2\omega'$. The condition (iii) is equivalent to asking that f_1 and g_1 be doubly periodic. One can see that the latter can be achieved by adjusting C and D if and only if

$$\begin{aligned} 2\mu k_1 + \sum_\alpha (u_\alpha + u'_\alpha) &\in 2\Omega, \\ 2\mu k_1 - \sum_\alpha (v_\alpha + v'_\alpha) &\in 2\Omega. \end{aligned} \quad (28)$$

Recalling that k_1 is real and positive, we conclude that there exist integers m, m', p, p' and a real number $x \in (0, 2\omega]$ such that $\sum_\alpha (u_\alpha + u'_\alpha) = -x +$

³There is a 4^m -fold ambiguity involved in the splitting $Q = Q_+ \cup Q_-$. It can be fixed, in principle, by requiring that the normal bundle of every section in the family be $\mathcal{O}(1) \oplus \mathcal{O}(1)$.

$2m\omega + 2m'\omega', \sum_{\alpha}(v_{\alpha} + v'_{\alpha}) = x + 2p\omega + 2p'\omega'$. Then Eqs. (28) together with the condition (iv) imply

$$2\mu k_1 = x. \quad (29)$$

Then for f_1 and g_1 to be doubly periodic one has to set

$$C = 2m\zeta_W(\omega) + 2m'\zeta_W(\omega'), D = 2p\zeta_W(\omega) + 2p'\zeta_W(\omega')$$

Let us notice for future use that

$$\begin{aligned} \log f_1(u + \omega) - \log f_1(u) &= -2\pi i m', \\ \log f_1(u + \omega') - \log f_1(u) &= 2\pi i m, \\ \log g_1(u + \omega) - \log g_1(u) &= -2\pi i p', \\ \log g_1(u + \omega') - \log g_1(u) &= 2\pi i p. \end{aligned} \quad (30)$$

Eq. (29) is a transcendental equation on k_1, k_2 , and the $SO(3)$ rotation required to bring S to the standard form Eq. (25). It reduces the number of real parameters in the equation of the curve from 5 to 4. Thus we have a four-parameter family of real sections of Z_2^0 .

5.3 The Kähler potential of the centered $n = 2$ moduli space

Having found a four-parameter family of real holomorphic sections of Z_2^0 we now would like to compute the corresponding hyperkähler metric. Since Z_2^0 has an intermediate holomorphic projection on $\mathcal{O}(4)$, we can use the method of Ref. [9] to write down the Legendre transform of the Kähler potential. The existence of the projection is equivalent to saying that η_2 is a holomorphic coordinate on Z_2^0 . The holomorphic 2-form ω in the patch $\zeta \neq \infty$ can be written as

$$\omega = d\eta_2 \wedge d \sum_{\text{branches}} \frac{1}{\eta} \log \frac{f_1}{g_1} \equiv d\eta_2 \wedge d\chi.$$

Here f_1 and $\eta = \sqrt{-\eta_2}$ are regarded as double-valued functions of $\zeta \in \mathbf{P}^1 \setminus \{\zeta \neq \infty\}$, and the sum is over the two branches of the cover $S \rightarrow \mathbf{P}^1$. Similarly, in the patch $\zeta \neq 0$ we can write

$$\omega' = d\eta'_2 \wedge d\chi'.$$

On the overlap we have the relations

$$\omega' = \zeta^{-2}\omega, \quad \eta'_2 = \zeta^{-4}\eta_2, \quad \chi' = \zeta^2\chi - 4\mu\zeta. \quad (31)$$

Following Ref. [9], we would like to find a (multi-valued) function $\hat{f}(\eta, \zeta)$ and a contour C on the double cover $S \rightarrow \mathbf{P}^1$ such that

$$\oint_C \frac{d\zeta}{\zeta^j} \hat{f}(\eta, \zeta) = \oint_0 \frac{d\zeta}{\zeta^{j-2}} \chi + \oint_\infty \frac{d\zeta}{\zeta^j} \chi'$$

for any integer j . Here the contours of integration on the RHS are small positively oriented loops around $\zeta = 0$ and $\zeta = \infty$. To find $\hat{f}$ we substitute the explicit expressions for χ and χ' and rewrite the integral on the RHS as an integral in the u -plane. Then the RHS becomes

$$\oint \frac{du}{k_1} \zeta(u)^{-j+2} \log \frac{f_1(u)}{g_1(u)} + 4\mu \oint_0 \frac{d\zeta}{\zeta^{j-1}}, \quad (32)$$

where the contour in the first integral consists of four small positively oriented loops around four preimages of the points $\zeta = 0$ and $\zeta = \infty$ in the fundamental rectangle of the lattice 2Ω . We denote these points $u_0, u'_0 = 2(\omega + \omega') - u_0, u_\infty, u'_\infty = 2(\omega + \omega') - u_\infty$. Besides these four points the only other branch points of $\log f_1(u)/g_1(u)$ in the fundamental rectangle are $u_\alpha, u'_\alpha, v_\alpha, v'_\alpha, \alpha = 1, \dots, k$. As for $\zeta(u)$, it is elliptic. Then we can rewrite Eq. (32) as

$$\oint_{bdry} \frac{du}{k_1} \zeta(u)^{-j+2} \log \frac{f_1(u)}{g_1(u)} + \sum_\alpha \oint_{A_\alpha + A'_\alpha} \frac{du}{k_1} \zeta(u)^{-j+2} \log \frac{f_1(u)}{g_1(u)} + 4\mu \oint_0 \frac{d\zeta}{\zeta^{j-1}}, \quad (33)$$

where the contour in the first integral runs along the boundary of the fundamental rectangle, while A_α and A'_α enclose the pairs of points u_α, v_α and u'_α, v'_α , respectively (see Figure 2). Using Eqs. (30) the integral over the boundary can be simplified to

$$-2\pi i \oint_{(m-p, m'-p')} \frac{du}{k_1} \zeta(u)^{-j+2},$$

where the contour $(m-p, m'-p')$ winds $m-p$ times around the real cycle and $m'-p'$ times around the imaginary cycle. Recalling the explicit form of $f_1(u)$ and $g_1(u)$, we can rewrite the integral over $A_\alpha + A'_\alpha$ as

$$\oint_{B_\alpha + B'_\alpha} \frac{du}{k_1} \zeta(u)^{-j+2} \log \sigma(u - u_\alpha) \sigma(u - u'_\alpha) \sigma(u - v_\alpha) \sigma(u - v'_\alpha), \quad (34)$$




Figure 2: Integration contours in Eq. (33). Only one of the contours A_α and one of the contours A'_α are shown.

where the contours B_α and B'_α are figure-eight-shaped contours shown in Figure 3. On the other hand, it can be easily seen that

$$\eta(u) - P_\alpha(\zeta(u)) \sim e^{u(C+D)} \frac{\sigma(u - u_\alpha)\sigma(u - u'_\alpha)\sigma(u - v_\alpha)\sigma(u - v'_\alpha)}{\sigma(u - u_\infty)^2\sigma(u + u_\infty)^2}.$$

Since neither u_∞ nor u'_∞ are enclosed by the contour $B_\alpha + B'_\alpha$, the integral Eq. (34) is equal to

$$\oint_{B_\alpha+B'_\alpha} \frac{du}{k_1} \zeta(u)^{-j+2} \log(\eta(u) - P_\alpha(\zeta(u))).$$

Collecting all of this together we get

$$\begin{aligned} \oint_C \frac{d\zeta}{\zeta^j} \hat{f}(\eta, \zeta) &= -2\pi i \oint_{(m-p, m'-p')} \frac{d\zeta}{\eta} \zeta^{-j+2} \\ &+ \sum_\alpha \oint_{C_\alpha+C'_\alpha} \frac{d\zeta}{\eta} \zeta^{-j+2} \log(\eta - P_\alpha(\zeta)) + 4\mu \oint_0 \frac{d\zeta}{\zeta^{j-1}}. \end{aligned} \quad (35)$$




Figure 3: The contours B_α and B'_α .

Here all the functions are regarded as functions on the double cover of the ζ -plane, and the contours C_α, C'_α are the images of B_α, B'_α under the map $u \mapsto \zeta$. We now define a function $G(\eta, \zeta)$ by $\partial G / \partial \eta = -2\eta\zeta^{-2}\hat{f}$. According to Ref. [9] the Legendre transform of the Kähler potential is given by

$$F = \frac{1}{2\pi i} \oint \frac{d\zeta}{\zeta^2} G(\eta, \zeta).$$

Hence we can read off F :

$$\begin{aligned} F = & -\frac{1}{2\pi i} \oint_0 d\zeta \frac{4\mu\eta^2}{\zeta^3} + \oint_{(m-p, m'-p')} d\zeta \frac{2\eta}{\zeta^2} \\ & - \sum_\alpha \frac{1}{2\pi i} \oint_{C_\alpha + C'_\alpha} \frac{d\zeta}{\zeta^2} 2(\eta - P_\alpha(\zeta)) \log(\eta - P_\alpha(\zeta)). \end{aligned} \quad (36)$$

F may be regarded as a function of the coefficients of $\eta_2(\zeta) = z + v\zeta + w\zeta^2 - \bar{v}\zeta^3 + \bar{z}\zeta^4$. Since w is real, F depends on 5 real parameters. These parameters are subject to one transcendental constraint expressed by Eq. (29). (This constraint implies $\partial F / \partial w = 0$.) Thus we may think of w as an implicit function

of z and v . The Kähler potential $K(z, \bar{z}, u, \bar{u})$ is the Legendre transform of F :

$$K(z, \bar{z}, u, \bar{u}) = F(z, \bar{z}, v, \bar{v}, w) - uv - \bar{u}\bar{v}, \quad \frac{\partial F}{\partial v} = u, \quad \frac{\partial F}{\partial \bar{v}} = \bar{u}.$$

Eq. (36) agrees with a conjecture by Chalmers [16].

We already saw in section 5 that M_2^0 is a resolution of D_k singularity. Now we can check that it is ALF. To this end we take the limit $k_1 \rightarrow +\infty$. Eq. (29) implies that in this limit $\omega \rightarrow \infty$, while ω' stays finite. Thus the curve S degenerates: $\eta_2(\zeta) \rightarrow -(P(\zeta))^2$, where $P(\zeta)$ is a real section of $\mathbf{T}$. It is easy to see that in this limit F reduces to the Taub-NUT form (see section 4):

$$F \sim \frac{1}{2\pi i} \oint_0 d\zeta \left(-\frac{4\mu P(\zeta)^2}{\zeta^3} + K \frac{P(\zeta) \log P(\zeta)}{\zeta^2} \right), \quad (37)$$

where K is an integer depending on the limiting behavior of $u_\alpha, u'_\alpha, v_\alpha, v'_\alpha$. Therefore asymptotically the metric on M_2^0 has the Taub-NUT form. With some more work it should be possible to compute the integer K as well.

Note also that if we set $\mu = 0$, then the metric becomes ALE. Kronheimer proved [2] that the D_k ALE metric is essentially unique. Thus we have obtained the Legendre transform of the Kähler potential for the D_k metrics of Ref. [1]. It would be interesting to obtain a similar representation for the E_k ALE metrics.

References

- [1] P.B. Kronheimer, “The Construction of ALE Spaces as Hyper-Kähler Quotients,” J. Differential Geometry **29**, 665-683 (1989).
- [2] P.B. Kronheimer, “A Torelli-type theorem for gravitational instantons,” J. Diff. Geom. **29** 685-697 (1989).
- [3] G.W. Gibbons and S.W. Hawking, “Gravitational Multi-instantons,” Phys. Lett. **B 78**, 430-432 (1978).
- [4] A. Sen, “A Note on Enhanced Gauge Symmetries in M- and String Theory,” JHEP **09**, 1 (1997) hep-th/9707123.
- [5] S.A. Cherkis and A. Kapustin, “ D_k Gravitational Instantons and Nahm Equations”, hep-th/9803112.

- [6] P.B. Kronheimer, “Monopoles and Taub-NUT Metrics,” M. Sc. Thesis, Oxford, 1985.
- [7] S.A. Cherkis and A. Kapustin, “Singular Monopoles and Supersymmetric Gauge Theories in Three Dimensions,” hep-th/9711145.
- [8] U. Lindström and M. Roček, Comm. Math. Phys. **115** 21 (1988).
- [9] I.T. Ivanov and M. Roček, “Supersymmetric Sigma Models, Twistors, and the Atiyah-Hitchin Metric,” Comm. Math. Phys. **182** 291-302 (1996).
- [10] N.J. Hitchin, “Monopoles and Geodesics,” Comm. Math. Phys. **83** 579-602 (1982).
- [11] J. Hurtubise and M. K. Murray, “On the Construction of Monopoles for the Classical Groups,” Comm. Math. Phys. **122** 35-89 (1989).
- [12] N.J. Hitchin, “On the Construction of Monopoles,” Comm. Math. Phys. **89** 145-190 (1983).
- [13] M. Atiyah and N. Hitchin, The Geometry and Dynamics of Magnetic Monopoles, Princeton Univ. Press, Princeton (1988).
- [14] N. J. Hitchin, A. Karlhede, U. Lindström, and M. Roček, “Hyperkähler Metrics and Supersymmetry,” Comm. Math. Phys. **108** 535-589 (1987).
- [15] W. Nahm, “Self-dual monopoles and calorons.” In: Lecture Notes in Physics **201**, G. Denardo et al. (eds.), Springer (1984).
- [16] G. Chalmers, “The Implicit Metric on a Deformation of the Atiyah-Hitchin Manifold,” hep-th/9709082, “Multi-monopole Moduli Spaces for $SU(N)$ Gauge Group,” hep-th/9605182.